

Antihistamines May Improve Response to Immunotherapy

Allergy meds were linked to longer survival among cancer patients treated with immune checkpoint inhibitors.

January 27, 2022 By [Sukanya Charuchandra](#)

Antihistamines, often used to treat allergies, are associated with better response to checkpoint inhibitor [immunotherapy](#) and improved overall survival, according to study results published in [Cancer Cell](#).

“In searching for factors that might influence responses to immunotherapy, we were surprised to discover that antihistamines, a mediator of the allergy response, were associated with significantly improved outcomes in patients,” Dihua Yu, MD, PhD, of the University of Texas MD Anderson Cancer Center, said in a [press release](#).

Checkpoint inhibitors block proteins on T cells that help regulate immune function. For example, some tumors can hijack a protein called PD-1 to turn off immune responses. Drugs that block the interaction between PD-1 on T cells and its binding partner on malignant cells can release the brakes and restore T-cell activity against cancer. Another checkpoint protein, CTLA-4, suppresses T-cell multiplication.

But not every individual with cancer responds to checkpoint inhibitor treatment, and the reasons are not fully understood. Yu and colleagues conducted a retrospective study to understand how antihistamines may boost antitumor immunity and alter response to immunotherapy.

The researchers conducted a retrospective study using data from people treated with immunotherapy at MD Anderson. Of 40 common drugs, H1-antihistamines alone were linked to improved survival. H1-antihistamines block the histamine receptor H1 (HRH1), preventing histamine—a chemical released by certain immune cells during allergic reactions—from interacting with its receptor.

They found that cancer patients with low plasma histamine levels had a higher overall response rate to PD-1 checkpoint inhibitors compared with patients who had high histamine levels. Among people who received checkpoint immunotherapy for melanoma or lung cancer, those who took antihistamines had significantly lower death rates and longer survival than those who did not take the common drugs. Although people with breast or colon cancer who took antihistamines also had

a reduced death rate, this association was not statistically significant. Antihistamines afforded little benefit to cancer patients treated with chemotherapy.

Data analyses of samples from [The Cancer Genome Atlas](#) revealed that higher levels of HRH1 in tumors were linked to poorer antitumor immunity, weaker response to immunotherapy and worse survival.

“Looking closer at this relationship, we discovered that histamine, through its receptor HRH1, can promote cancer cell immune evasion and resistance to immunotherapy,” study coauthor Yi Xiao, PhD, said in the press release.

Further, the researchers noted that both histamine and HRH1 were present at high levels within the tumor microenvironment. The HRH1 receptor was highly expressed on certain types of macrophages that suppress immune response, and histamine was present in cancer cells. When they blocked HRH1, either by knocking out genes or use of antihistamines, immunosuppression was reversed, resulting in increased T-cell activation that checked tumor growth.

In addition, the researchers used mouse models to test the effect of allergic reactions and antihistamines on immunotherapy response and tumor growth. While histamine levels increased in parallel with tumor growth, this could be countered with antihistamines. In a different set of experiments, they discovered a link between plasma histamine levels and immunotherapy response in people with cancer. Taken together, these findings suggest that pre-existing allergies or high histamine levels can lead to tamped-down immunotherapy response.

“Our preclinical findings suggest that antihistamines have the potential to enhance responses to immunotherapy, especially in those with high levels of histamine in the blood,” said Yu. “There is more work to be done, but we are excited to continue exploring possible therapeutic applications with antihistamines, which offer an inexpensive approach with minimal side effects.”

The researchers now plan to design prospective clinical trials to explore antihistamines as part of combination regimens with checkpoint inhibitor immunotherapy.

Click here to read the study abstract in [Cancer Cell](#).

Click here to learn more about [immunotherapy for cancer](#)