

HCV-Positive and HCV-Negative Kidney Transplants Have Similar Outcomes

Study findings suggest HCV-RNA-positive kidneys are not inferior in the direct-acting antiviral era.

November 16, 2022 By [Sukanya Charuchandra](#)

People who receive kidney transplants from deceased donors with hepatitis C virus (HCV) have graft survival rates comparable to those who receive HCV-negative organs, according to study findings published in [JAMA](#).

“This study and others suggest that many patients on the transplant waiting list should weigh the option of transplant with an HCV-RNA-positive donor kidney,” the study authors wrote.

Many deceased organ donors in the United States have [hepatitis C](#). Before the advent of [direct-acting antiviral \(DAA\) therapy](#), recipients of organs from donors with HCV were at risk of developing the disease themselves, which could lead to transplant failure and even death. Many HCV-positive organs were therefore discarded. But DAAs, which can cure hepatitis C in nearly all patients, have made transplants from HCV-positive donors safe, with trials showing 100% cure rates.

The Kidney Donor Profile Index (KDPI) is a measure of the quality ascribed to donor kidneys, with higher values suggesting poorer quality. Based on data from the period when older interferon-based therapy was used to treat hepatitis C, the index considers kidneys from HCV-positive donors to be inferior, and a high KDPI score may be one reason why such kidneys continue to be discarded.

Peter Reese, MD, PhD, of the Perelman School of Medicine at the University of Pennsylvania, and colleagues assessed whether the penalty levied on kidneys from HCV-RNA-positive donors is relevant when effective antivirals are available.

The researchers carried out a retrospective cohort study of adults who received kidney transplants from deceased donors between July 2016 and December 2021 using data from the Organ Procurement and Transplantation Network.

The study population included 45,827 deceased donors and 75,905 kidney recipients at 217 U.S. centers. Ultimately, the team compared outcomes for recipients of 2,551 kidneys from HCV-RNA-positive donors and 43,276 organs from HCV-RNA-negative donors, looking at five-year kidney

graft survival rates. They focused on HCV RNA status (detectable or undetectable), which indicates active infection; recipients of kidneys that were HCV antibody positive but HCV RNA negative were excluded.

Deceased donors with hepatitis C were younger, on average, than their HCV-negative counterparts (median 35 versus 39 years), and they were less likely to have hypertension or diabetes or to have suffered a stroke. Additionally, their donated kidneys had better KDPI scores (32% versus 47%). Recipients of HCV-positive kidneys were older, on average (60 versus 56 years), weighed more and were more likely to have diabetes than recipients of HCV-negative organs.

The researchers saw little difference in five-year graft survival rates for people who received kidneys from HCV-RNA-positive versus HCV-RNA-negative donors. Those who received transplants from donors with HCV had a 72% five-year graft survival rate compared with 69% for those who received kidneys from donors without HCV. The average survival time for the kidney graft was 4.30 years for those with HCV-positive transplants versus 4.27 years for those who received HCV-negative organs.

“Recipients of kidney transplants from HCV-RNA-positive deceased donors had five-year mean allograft survival that was not statistically different from HCV-RNA-negative kidney recipients during the direct-acting antiviral era,” wrote the researchers. “These findings may provide a rationale for the transplant community to reexamine the KDPI’s HCV penalty.”

Click here to read the study in [JAMA](#).

Click here for [more news about hepatitis C](#).